



National
Comprehensive
Cancer
Network®

NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®)

AIDS-Related Kaposi Sarcoma

Version 1.2018 — November 3, 2017

NCCN.org

Continue



National
Comprehensive
Cancer
Network®

NCCN Guidelines Version 1.2018 Panel Members

AIDS-Related Kaposi Sarcoma

[NCCN Guidelines Index](#)
[Table of Contents](#)
[Discussion](#)

* Gita Suneja, MD/Co-chair §
Duke Cancer Institute

* Erin Reid, MD/Co-chair ‡
UC San Diego Moores Cancer Center

Richard F. Ambinder, MD, PhD †
The Sidney Kimmel Comprehensive
Cancer Center at Johns Hopkins

* Kevin Ard, MD, MPH ☐ ♯
Massachusetts General Hospital
Cancer Center

Robert Baiocchi, MD, PhD †
The Ohio State University Comprehensive
Cancer Center - James Cancer Hospital
and Solove Research Institute

Stefan K. Barta, MD, MRCP † ‡ ♯
Fox Chase Cancer Center

Evie Carchman, MD ¶
University of Wisconsin
Carbone Cancer Center

Adam Cohen, MD †
Huntsman Cancer Institute
at the University of Utah

Neel Gupta, MD †
Stanford Cancer Institute

Kimberly L. Johung, MD, PhD §
Yale Cancer Center/Smilow Cancer Hospital

Ann Klopp, MD, PhD §
The University of Texas
MD Anderson Cancer Center

Ann S. LaCasce, MD †
Dana-Farber/Brigham and Women's
Cancer Center

Chi Lin, MD, PhD §
Fred & Pamela Buffett Cancer Center

Oxana V. Makarova-Rusher, MD †
University of Michigan
Comprehensive Cancer Center

Amitkumar Mehta, MD ‡
University of Alabama at Birmingham
Comprehensive Cancer Center

* Manoj P. Menon, MD, MPH †
Fred Hutchinson Cancer Research Center/
Seattle Cancer Care Alliance

David Morgan, MD ‡
Vanderbilt-Ingram Cancer Center

Nitya Nathwani, MD ‡
City of Hope Comprehensive Cancer Center

* Ariela Noy, MD ‡
Memorial Sloan Kettering Cancer Center

Frank Palella, MD
Robert H. Lurie Comprehensive Cancer
Center of Northwestern University

Lee Ratner, MD, PhD † ♯
Siteman Cancer Center at Barnes-Jewish Hospital
and Washington University School of Medicine

Stacey Rizza, MD ☐
Mayo Clinic Cancer Center

Michelle A. Rudek, PhD, PharmD ∑
The Sidney Kimmel Comprehensive
Cancer Center at Johns Hopkins

Jeff Taylor ¥
Coachella Valley Community Research Initiative

Benjamin Tomlinson, MD † ‡
Case Comprehensive Cancer Center/
University Hospitals Seidman Cancer Center and
Cleveland Clinic Taussig Cancer Institute

* Chia-Ching J. Wang, MD †
UCSF Helen Diller Family
Comprehensive Cancer Center

NCCN
Mary Dwyer, MS
Deborah Freedman-Cass, PhD

Continue

[NCCN Guidelines Panel Disclosures](#)

† Medical oncology	♯ Internal medicine
‡ Hematology/Hematology oncology	∑ Pharmacology/Pharmacy
§ Radiotherapy/Radiation oncology	☐ Infectious diseases
¶ Surgery/Surgical oncology	¥ Patient advocacy
	* Discussion Writing Committee Member



[NCCN AIDS-Related Kaposi Sarcoma Panel Members](#)

- [Diagnosis and Workup \(KS-1\)](#)
- [Limited Cutaneous Disease \(KS-2\)](#)
- [Advanced Cutaneous, Oral, Visceral, or Nodal Disease \(KS-3\)](#)
- [Surveillance \(KS-4\)](#)

- [Staging Classification for AIDS-Related KS \(KS-A\)](#)
- [Principles and Goals of Therapy \(KS-B\)](#)
- [Local Therapy \(KS-C\)](#)
- [Systemic Therapy \(KS-D\)](#)

Clinical Trials: NCCN believes that the best management for any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

To find clinical trials online at NCCN Member Institutions, [click here: nccn.org/clinical_trials/physician.html](#).

NCCN Categories of Evidence and Consensus: All recommendations are category 2A unless otherwise indicated.

See [NCCN Categories of Evidence and Consensus](#).

The NCCN Guidelines® are a statement of evidence and consensus of the authors regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult the NCCN Guidelines is expected to use independent medical judgment in the context of individual clinical circumstances to determine any patient's care or treatment. The National Comprehensive Cancer Network® (NCCN®) makes no representations or warranties of any kind regarding their content, use or application and disclaims any responsibility for their application or use in any way. The NCCN Guidelines are copyrighted by National Comprehensive Cancer Network®. All rights reserved. The NCCN Guidelines and the illustrations herein may not be reproduced in any form without the express written permission of NCCN. ©2017.



NCCN Guidelines Version 1.2018

AIDS-Related Kaposi Sarcoma

DIAGNOSIS

ESSENTIAL:

- Review of all slides with at least one paraffin block representative of the tumor by a pathologist with expertise in the diagnosis of Kaposi sarcoma (KS).
 - Rebiopsy if nondiagnostic
- Histopathology review of adequate biopsy (ie, skin punch, incisional, excisional)
- Adequate immunophenotyping to establish diagnosis
- IHC panel: KSHV (HHV-8) LANA-1

USEFUL IN CERTAIN CIRCUMSTANCES:

- IHC: CD31 and CD34 if unclear whether tumor has a vascular origin
- Encourage additional biopsy of nodal or visceral sites, if a coexisting disorder is suspected (ie, infection, lymphoma, multicentric Castleman's disease)

WORKUP

ESSENTIAL:

- History and physical exam
 - including history of additional immunosuppression such as transplant/glucocorticoids
 - including complete skin and oral exams and documentation of edema
- CBC, differential, comprehensive metabolic panel
- Quantitative HIV viral load^a
- T-cell subsets^a
- Evaluation for suspected opportunistic infections^a
- Stool hemocult
- HIV diagnostic testing, if not already performed
- Chest x-ray
- Photography of oral, conjunctival, and cutaneous lesions (with reference unit of measure in picture) for documentation of extent of disease
- Pregnancy testing in women of child-bearing age (if chemotherapy or RT planned)

USEFUL IN SELECTED CASES:

- Chest CT with contrast ± abdominal/pelvic CT with contrast or MRI with contrast and/or PET/CT scan^b
- Upper endoscopy (EGD)/colonoscopy if GI symptoms or positive hemocult
- Bronchoscopy if unexplained pulmonary symptoms or abnormalities on chest x-ray or CT/MRI
- Transthoracic echocardiogram, if anthracycline planned or suspected pericardial effusion
- Lab workup of coexisting HHV-8–associated diseases^c

AIDS-RELATED KS STAGE^d

Limited cutaneous disease

→ [See First-Line Therapy \(KS-2\)](#)

Advanced cutaneous, oral, visceral, or nodal disease

→ [See First-Line Therapy \(KS-3\)](#)

^aAll HIV seropositive patients should have recent T-cell subsets, including quantitative CD4+ T-cell count, and HIV viral load to assess immune function and HIV control ([see Discussion](#)). Involvement of an infectious disease (ID) specialist to evaluate for coexisting opportunistic infection (OI) is appropriate, especially with advanced immunosuppression.

^bImaging should be directed by symptoms or findings concerning for visceral or bone involvement as well as coexisting KSHV-associated inflammatory cytokine syndrome (KICS), multicentric Castleman's disease (MCD), or HHV8+ lymphoma; imaging is standard for staging of transplant-associated Kaposi sarcoma (KS).

^cUseful in setting of clinical features (ie, fever, dyspnea, effusions) concerning for KICS or KSHV-associated MCD: C-reactive protein, KSHV serum viral load, SPEP, IL-6, or IL-10.

^d[See Staging Classification for AIDS-Related KS \(KS-A\)](#).


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2018

AIDS-Related Kaposi Sarcoma



^eSee [Principles and Goals of Therapy \(KS-B\)](#).

^fInitiation of ART may result in immune reconstitution inflammatory syndrome (IRIS) within 3–6 months; IRIS is characterized by marked lesional swelling, increased tenderness, and peripheral edema. However, ART should not be delayed or discontinued unless life-threatening IRIS develops. Reconstitution of immune function is important for obtaining and maintaining control or remission of KS.

^gGlucocorticoids in any formulation should be avoided due to their association with KS progression. However, in cases of life-threatening conditions including IRIS, their use may be considered.

^hSee [clinicaltrials.gov](#).

ⁱSee [Local Therapy \(KS-C\)](#).

^jSee [Systemic Therapy \(KS-D\)](#).


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2018

AIDS-Related Kaposi Sarcoma



^eSee Principles and Goals of Therapy (KS-B).

^fInitiation of ART may result in IRIS within 3–6 months; IRIS is characterized by marked lesional swelling, increased tenderness, and peripheral edema. However, ART should not be delayed or discontinued unless life-threatening IRIS develops. Reconstitution of immune function is important for obtaining and maintaining control or remission of KS.

^gGlucocorticoids in any formulation should be avoided due to their association with KS progression. However, in cases of life-threatening conditions including IRIS, their use may be considered.

^hSee clinicaltrials.gov.

ⁱSee Local Therapy (KS-C).

^jSee Systemic Therapy (KS-D).

^kSystemic therapy is preferred over radiation therapy as first-line therapy and relapsed/refractory therapy for disseminated disease whenever systemic therapy is feasible considering performance status and comorbidities.

^lPatients with relapsed/refractory disease who received RT in first-line therapy should receive systemic therapy if possible.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



SURVEILLANCE

- **For patients not requiring active therapy and with no signs of progression**
 - ▶ **Every 3 months for year 1, then every 4-6 months for year 2, then every 6-12 months thereafter**
 - ◇ **History and physical exam**
 - including history of additional immunosuppression such as transplant/glucocorticoids
 - including complete skin and oral exams, and documentation of edema
 - ◇ **CBC, differential, comprehensive metabolic panel, T-cell subsets (CD4+ T-cell count), and HIV viral load**
 - ◇ **Assess ART compliance**
- **Photography of oral, conjunctival, and cutaneous lesions (with reference unit of measure in picture) for documentation of extent of disease if change in disease is noted**
- **If signs and symptoms concerning for visceral involvement or prior to new therapy if progression/refractory disease**
 - ▶ **Stool hemoccult**
 - ▶ **Chest x-ray or chest CT with contrast**
 - ▶ **EGD/colonoscopy**
 - ▶ **Bronchoscopy**
- **As KSHV is not eradicated with treatment of KS, the risk for future KS persists even after complete remission. Optimization and monitoring of HIV control and immune function is important to minimize this risk. This risk depends on immune function and generally decreases with immune reconstitution. However, KS can persist, relapse, or present even in the setting of normal values of T-cell subsets. Less frequent (every 6–12 mo) oncology monitoring may be appropriate for selected patients with undetectable HIV viral loads, normal T-cell subsets, and stable KS for 2 or more years as long as the patient has regular follow-up with an HIV provider.**

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2018

AIDS-Related Kaposi Sarcoma

STAGING CLASSIFICATION FOR AIDS-RELATED KS*

	Good risk (all of the following)	Poor risk (any of the following)
Tumor, T	T0: Confined to skin and/or lymph nodes and/or minimal oral disease (non-nodular KS confined to palate)	T1: Tumor-associated edema or ulceration Extensive oral KS Gastrointestinal KS KS in organs other than lymph nodes
Immune system, I¹	I0: CD4+ T-cell count $\geq 150/\mu\text{L}$	I1: CD4+ T-cell count $< 150/\mu\text{L}$
Systemic disease, S	S0: No history of opportunistic infection or thrush No “B” symptoms² Karnofsky performance status ≥ 70	S1: History of opportunistic infection and/or thrush “B” symptoms present Karnofsky performance status < 70 Other HIV-related illness (eg, neurologic disease, lymphoma)
¹ I stage has less prognostic value than T or S stages in the presence of ART therapy ² “B” symptoms are unexplained fever, night sweats, > 10 percent involuntary weight loss, or diarrhea persisting more than two weeks		

*Adapted from Krown SE, Metroka C, Wernz JC. Kaposi's sarcoma in the acquired immune deficiency syndrome: a proposal for uniform evaluation, response, and staging criteria. AIDS Clinical Trials Group Oncology Committee. J Clin Oncol 1989;7:1201-1207.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2018

AIDS-Related Kaposi Sarcoma

PRINCIPLES AND GOALS OF THERAPY

PRINCIPLES OF THERAPY:

- **Individual KS lesions may be distinct clones that arise due to the common risk factors of immunosuppression and persistent HHV-8 infection as opposed to metastases. Treatment of existing disease therefore may not prevent occurrence of future lesions.**
- **Reconstitution of immune function, maintenance of viral suppression, and avoidance of additional immunosuppression are critical to prevention of additional KS lesions and maintenance of response to therapy. For AIDS-related KS, it is important to work with an HIV specialist to optimize suppression of HIV and reconstitution of immune function with ART. Important examples of iatrogenic immunosuppression, which may promote KS, include not only systemic but local glucocorticoids (ie, inhaled, topical, intra-articular). Note that KS may flare in a remote location from the site of local glucocorticoids. Patients requiring rituximab for treatment of NHL with coexisting KS or multicentric Castleman's disease may develop flares of KS or incident KS. This may be mitigated by use of concurrent chemotherapy active against both KS and disease for which rituximab is prescribed (ie, doxorubicin).**
- **Persons with AIDS-related KS, especially those with advanced immunosuppression, are at increased risk of opportunistic infections (OIs), marrow suppression with neutropenic fever, or thrombocytopenic bleeding and should be monitored closely. It is important to collaborate with an HIV specialist to ensure adequate OI prophylaxis appropriate to CD4+ T-cell count (which may temporarily decrease with cytotoxic chemotherapy). Growth factor support may be needed to facilitate systemic therapy.**
- **Lymphedema and soft tissue infections: KS is often complicated by lymphedema with increased risk of cellulitis and deep tissue infections in affected limbs. Risk of severe lymphedema and delayed wound healing may be increased after radiation. Refer to a lymphedema specialist. In the setting of advanced cutaneous disease, radiation should be reserved for circumstances when systemic therapy is not feasible with the goal of palliation or short-term disease management until systemic therapy may be delivered. Note that treatment responses may be delayed in the context of significant lymphedema.**

GOALS OF THERAPY:

- **Patients with limited cutaneous disease that is asymptomatic and cosmetically acceptable may be observed while continuing ART with optimization of immune function and HIV viral suppression as above. Remissions or stable disease may occur with ART and optimization of immune function and HIV viral suppression alone.**
- **Patients with symptomatic or cosmetically unacceptable disease should use the most minimally invasive and least toxic therapy to control disease. A limited number of cycles of systemic therapy (eg, 3–6) may be sufficient for those initiating or re-initiating ART.**
- **Patients with advanced symptomatic cutaneous, visceral, nodal, or oral disease should be treated with systemic therapy with the goal of reducing or reversing symptoms, lymphedema, or threat to organ function. Complete remissions are rare.**
 - ▶ **Treatment is typically continued until unacceptable toxicity or plateau in response; maintenance therapy beyond 2 cycles of systemic therapy after determination of plateau is not recommended. If response is then clinically acceptable, patients may be observed on ART alone. Otherwise, alternative therapy should be initiated.**

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

LOCAL THERAPY**Topical**

- **Alitretinoin 0.1% gel¹**
 - ▶ **Apply 3–4 times daily to affected skin sites**
- **Imiquimod, 5% cream²**
 - ▶ **Apply 20 cm² of skin sachet under occlusion 3 times weekly; titrate dose to effect, tolerability**

Intralesional chemotherapy

- **Vinblastine³**
 - ▶ **0.2 mg/mL solution with a volume of 0.1 mL per 0.5 cm² of lesion**
 - ◇ **Other treatment schemas have been studied, with a variety of vinblastine concentrations, doses, administration volumes, frequency of administration, and total doses/volumes administered. See [Discussion](#) for additional references and information**
 - ▶ **Pain from injection is common and may persist for several days. NSAID may be useful to relieve pain from injection.**

Radiotherapy^{4,5,6}

- **24 Gy in 12 fractions in 2.0 Gy per fraction**
- **Other dosing schemas ranging from 6–8 Gy in 1 fraction to 30 Gy in 10–15 fractions may be used**
- **For most skin lesions, electrons or superficial x-rays can be used to deliver optimal dosimetry and minimize dose to underlying structures. To ensure sufficient dose is delivered for deeper or larger lesions, conformal photon therapy or mixed photon-electron treatment plans may be utilized. IMRT with or without image guidance may be useful for larger lesions. The use of bolus may be necessary to achieve adequate skin dose.**

¹Bodsworth NJ, Bloch M, Bower M, et al. Phase III vehicle-controlled, multi-centered study of topical alitretinoin gel 0.1% in cutaneous AIDS-related Kaposi's sarcoma. *Am J Clin Dermatol* 2001;2:77-87.

²Schatz NEC, Chevret S, Paz C, et al. Imiquimod 5% cream for treatment of HIV-negative Kaposi's sarcoma skin lesions: a phase I to II open-label trial. *J Am Acad Dermatol* 2008;58:585-591.

³Epstein JB. Treatment of oral Kaposi sarcoma with intralesional vinblastine. *Cancer* 1993;71:1722-1725.

⁴Singh NB, Lakier RH, Donde B. Hypofractionated radiation therapy in the treatment of epidemic Kaposi sarcoma – a prospective randomized trial. *Radiother Oncol* 2008;88:211-216.

⁵Hauerstock D1, Gerstein W, Vuong T. Results of radiation therapy for treatment of classic Kaposi sarcoma. *J Cutan Med Surg* 2009;13:18-21.

⁶Kirova YM1, Belembaogo E, Frikha H, et al. Radiotherapy in the management of epidemic Kaposi's sarcoma: a retrospective study of 643 cases. *Radiother Oncol* 1998;46:19-22.

Note: All recommendations are category 2A unless otherwise indicated.


Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 1.2018

AIDS-Related Kaposi Sarcoma

SYSTEMIC THERAPY^a



^aSee references for regimens on [KS-D 2 of 2](#).

^bDue to risk of cardiotoxicity, perform echocardiogram prior to initial and repeat course of liposomal doxorubicin and limit lifetime dose to 400–450 mg/m².

^cConsider repeating any prior systemic therapy that was tolerated and resulted in a durable response.

^dIf both first-line options have already been given, the patient should proceed to the subsequent systemic therapy options.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



SYSTEMIC THERAPY

REFERENCES

- ¹Northfelt DW, Dezube BJ, Thommes JA, et al. Pegylated-liposomal doxorubicin versus doxorubicin, bleomycin, and vincristine in the treatment of AIDS-related Kaposi's sarcoma: results of a randomized phase III clinical trial. *J Clin Oncol* 1998;16:2445-2451.
- ²Cianfrocca M, Lee S, Von Roenn J, et al. Randomized trial of paclitaxel versus pegylated liposomal doxorubicin for advanced human immunodeficiency virus-associated Kaposi sarcoma: evidence of symptom palliation from chemotherapy. *Cancer* 2010;116:3969-3977.
- ³Polizzotto MN, Uldrick TS, Kyvill KM, et al. Pomalidomide for symptomatic Kaposi's sarcoma in people with and without HIV infection: a phase I/II study. *J Clin Oncol* 2016;34:4125-4131.
- ⁴Uldrick TS, Wyvill KM, Kumar P, et al. Phase II study of bevacizumab in patients with HIV-associated Kaposi's sarcoma receiving antiretroviral therapy. *J Clin Oncol* 2012;30:1476-1483.
- ⁵Evans SR, Krown SE, Testa MA, et al. Phase II evaluation of low-dose oral etoposide for the treatment of relapsed or progressive AIDS-related Kaposi's sarcoma: an AIDS Clinical Trials Group clinical study. *J Clin Oncol* 2002;20:3236-3241.
- ⁶Strother RM, Gregory KM, Pastakia SD, et al. Retrospective analysis of the efficacy of gemcitabine for previously treated AIDS-associated Kaposi's sarcoma in western Kenya. *Oncology* 2010;78:5-11.
- ⁷Koon HB, Krown SE, Lee JY, et al. Phase II trial of imatinib in AIDS-associated Kaposi's sarcoma: AIDS Malignancy Consortium Protocol 042. *J Clin Oncol* 2014;32:402-408.
- ⁸Krown SE, Li P, Von Roenn JH, et al. Efficacy of low-dose interferon with antiretroviral therapy in Kaposi's sarcoma: a randomized phase II AIDS clinical trials group study. *J Interferon Cytokine Res* 2002;22:295-303.
- ⁹Fortino S, Santoro M, Iuliano E, et al. Treatment of Kaposi's sarcoma (KS) with nab-paclitaxel. *Ann Oncol* 2016;27:iv124.
- ¹⁰Little RF, Wyvill KM, Pluda JM, et al. Activity of thalidomide in AIDS-related Kaposi's sarcoma. *J Clin Oncol* 2000;18:2593-2602.
- ¹¹Nasti G, Errante D, Talamini R, et al. Vinorelbine is an effective and safe drug for AIDS-related Kaposi's sarcoma: results of a phase II study. *J Clin Oncol* 2000;18:1550-1557.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



Discussion

NCCN Categories of Evidence and Consensus

Category 1: Based upon high-level evidence, there is uniform NCCN consensus that the intervention is appropriate.

Category 2A: Based upon lower-level evidence, there is uniform NCCN consensus that the intervention is appropriate.

Category 2B: Based upon lower-level evidence, there is NCCN consensus that the intervention is appropriate.

Category 3: Based upon any level of evidence, there is major NCCN disagreement that the intervention is appropriate.

All recommendations are category 2A unless otherwise noted.

Local Excision.....	MS-8
Radiation Therapy	MS-8
Systemic Therapy	MS-8
Surveillance of Patients with AIDS-Related Kaposi Sarcoma ...	MS-9
Systemic Therapy of Relapsed/Refractory Disease.....	MS-10
Summary.....	MS-11
References.....	MS-13

Table of Contents

Overview	MS-2
Literature Search Criteria and Guidelines Update Methodology ...	MS-2
HIV Management during Cancer Therapy	MS-3
HIV Screening.....	MS-3
Linkage to HIV Care.....	MS-3
HIV Therapy during Cancer Treatment	MS-3
Opportunistic Infection Prophylaxis.....	MS-4
AIDS-Related Kaposi Sarcoma	MS-4
Diagnosis and Workup of AIDS-Related Kaposi Sarcoma	MS-5
Staging of AIDS-Related Kaposi Sarcoma	MS-6
Initial Management of AIDS-Related Kaposi Sarcoma	MS-6
Antiretroviral Therapy (ART)	MS-7
Topical Therapies	MS-7
Intralesional Chemotherapy	MS-7

Overview

In 2017, it is estimated that more than 1.1 million people in the United States are living with human immunodeficiency virus (HIV) infection.¹ Without treatment, HIV infection causes the Acquired Immune Deficiency Syndrome (AIDS) and AIDS-defining cancers: non-Hodgkin's lymphoma, Kaposi sarcoma, and cervical cancer.^{2,3} Dramatically improved treatment of HIV over the last two decades has decreased the risk of AIDS, improved immune function and survival, and reduced AIDS-defining cancers in this population.^{4,5} As people living with HIV (PLWH) live longer and healthier lives, however, they experience an increased risk of many non-AIDS-defining cancers.⁶⁻¹⁰

It is estimated that 7760 PLWH were diagnosed with cancer in the United States in 2010, representing an approximately 50% increase over the expected number in the general population.¹¹ Other studies have also noted a higher risk for developing cancer in PLWH than in HIV-negative individuals, likely due to underlying immune dysregulation and co-infection with viruses such as human papillomavirus (HPV), human herpesvirus 8 (HHV-8), hepatitis B virus (HBV), hepatitis C virus (HCV), and Epstein Barr virus (EBV).¹²⁻¹⁶ In addition, the prevalence of other cancer risk factors in the HIV-positive population (eg, smoking, heavy alcohol consumption) may play a role.¹⁷⁻²¹

The proportions of each major cancer type among total incident cancer cases occurring in PLWH in the U.S. during 2010 are:¹¹

- Non-Hodgkin's lymphoma 21%
- Kaposi sarcoma 12%
- Lung cancer 11%
- Anal cancer 10%
- Prostate cancer 7%
- Liver cancer 5%

- Colorectal cancer 5%
- Hodgkin lymphoma 4%
- Oral/pharyngeal cancer 4%
- Female breast cancer 2%
- Cervical cancer 1%

The NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for AIDS-Related Kaposi Sarcoma provide treatment recommendations for HIV seropositive individuals who develop Kaposi sarcoma and are intended to assist healthcare providers with clinical decision-making. This Discussion section provides an overview of the literature supporting the recommendations included in the guidelines. The panel will soon also publish separate NCCN Guidelines for Cancer in People Living With HIV, which will give recommendations for the management of non-small cell lung cancer (NSCLC), anal cancer, Hodgkin lymphoma, and cervical cancer in PLWH. Those guidelines will also offer general advice for this population regarding HIV management during cancer therapy, drug-drug interactions with antiretrovirals and cancer therapies, radiation therapy, and supportive care. Recommendations for the management of Non-Hodgkin's lymphoma in PLWH are available in the NCCN Guidelines for B-cell Lymphomas (available at www.NCCN.org).

Literature Search Criteria and Guidelines Update Methodology

Prior to the development of the NCCN Guidelines for Cancer in People Living with HIV, an electronic search of the PubMed database was performed to obtain key literature in the field published between April 11, 2007 and April 11, 2017, using the following search terms: (cancer or malignancy or carcinoma or adenocarcinoma or lymphoma or leukemia or melanoma or sarcoma or neoplasia) and (HIV or AIDS).



The PubMed database was chosen because it remains the most widely used resource for medical literature and indexes only peer-reviewed biomedical literature.⁶

The search results were narrowed by selecting studies in humans published in English. Results were confined to the following article types: Clinical Trial, Phase II; Clinical Trial, Phase III; Clinical Trial, Phase IV; Practice Guideline; Randomized Controlled Trial; Meta-Analysis; Systematic Reviews; and Validation Studies.

The PubMed search resulted in 771 citations, and their potential relevance was examined. The data from key PubMed articles and articles from additional sources deemed as relevant to these Guidelines and discussed by the panel have been included in this version of the Discussion section (eg, e-publications ahead of print, meeting abstracts). Recommendations for which high-level evidence is lacking are based on the panel's review of lower-level evidence and expert opinion.

The complete details of the Development and Update of the NCCN Guidelines are available on the NCCN website (www.NCCN.org).

HIV Management during Cancer Therapy

HIV Screening

One out of every 7 people in the United States who are infected with HIV (or approximately 157,000 people) are not aware of their infection status.¹ Infected individuals who are unaware of their HIV status do not receive the clinical care they need to reduce HIV-related morbidity and mortality and may unknowingly transmit HIV.²² The Centers for Disease Control and Prevention (CDC) therefore recommends HIV screening for all patients in all healthcare settings unless the patient declines testing (opt-out screening).²³

HIV testing may be particularly important in patients with cancer, because identification of HIV infection has the potential to improve clinical outcomes.²⁴ Results of a retrospective cohort study at MD Anderson Cancer Center revealed, however, that the rate of HIV testing from 2007 to 2009 was only 19.3%.²⁵ Analysis of data from the 2009 Behavioral Risk Factor Surveillance System showed that 41% of U.S. cancer survivors <65 years of age reported ever being tested for HIV.²⁶ In both studies, race and other demographic characteristics as well as tumor type influenced the likelihood of receiving an HIV test.

The NCCN panel supports the CDC recommendation and believes that all patients diagnosed with cancer who do not opt-out should be tested for HIV if their HIV status is unknown.

Linkage to HIV Care

The HIV Care Continuum Initiative indicates that all patients diagnosed with HIV should be connected with an HIV specialist.²⁷ Linkage to care with an HIV specialist has been shown to improve viral suppression and care engagement.^{28,29} Patients should initiate and continue antiretroviral therapy (ART) to achieve and maintain viral suppression and immune reconstitution. Early initiation of ART has been shown to improve survival in PLWH.³⁰ Linkage to HIV care is also essential for PLWH who have cancer, and the oncology team should refer all PLWH who have cancer to an HIV specialist, if they are not already linked to one. In all cases, communication between the oncologist and HIV specialist should be established. The HIV.gov website has a map that can be used to locate HIV services: <https://locator.hiv.gov/>.

HIV Therapy during Cancer Treatment

If the patient has already started ART, it should be continued during cancer treatment. For patients who have not yet started antiviral treatment, ART should optimally be initiated ≥ 7 days prior to start of

cancer treatment or after the first cycle of cancer therapy to facilitate separate assessment of tolerability of ART and cancer treatment.

ART interruptions during cancer treatment should generally be avoided, because they increase the risk of immunologic compromise, opportunistic infection, and death.³¹ Continuation of ART also may result in better tolerance of cancer treatment, higher response rates, and improved survival.^{32,33} If drug-drug interactions between cancer treatment and ART are problematic, then alternative ART regimens can be used.

Laboratory testing, including HIV viral load and CD4+ T cell monitoring, should generally be performed as per normal schedules in conjunction with the patient's HIV specialist.²¹ However, more frequent HIV viral load testing (eg, once a month for the first 3 months and then every 3 months) may be needed if systemic cancer therapy is used.³⁴

Opportunistic Infection Prophylaxis

The occurrence of opportunistic infections in PLWH has decreased in the ART era, mainly because effective ART reduces infection risk as CD4+ T cell counts rise.^{5,35-37} Furthermore, prophylaxis and treatment of opportunistic infections in PLWH have improved.^{37,38} Still, opportunistic infections represent a major cause of morbidity and mortality in PLWH.^{37,38}

The risk of bacterial, fungal, and viral infections is also elevated in patients with cancer, who may experience immunosuppression resulting from cancer treatment and sometimes from the disease itself (eg, hypogammaglobulinemia in lymphoid malignancies).³⁹⁻⁴³ In particular, chemotherapy can cause neutropenia, which is a major risk factor for the development of infections.⁴⁴ Newer targeted agents are also associated with immunosuppression and increased infection risk.⁴⁵ The

frequency and severity of infection are inversely proportional to the neutrophil count, with the risks of severe infection and bloodstream infection greatest (approximately 10% to 20%) at neutrophil counts below 100 cells/mcL.⁴⁶

PLWH may be more susceptible to infectious complications following chemotherapy than their uninfected counterparts, and low CD4+ T cell counts appear to increase the risk of febrile neutropenia.⁴⁷ Furthermore, data show that certain chemotherapy regimens can cause a sustained drop in CD4+ T cell counts and an increased risk of opportunistic infections.⁴⁸ Other regimens, however, appear to have similar effects on myelosuppression and infectious complications in PLWH who have cancer and HIV-negative patients with cancer.⁴⁹

Overall, the NCCN panel believes that PLWH who have cancer should receive the prophylaxis indicated by their HIV status, as recommended in U.S. Department of Health and Human Services' Guidelines for the Prevention and Treatment of Opportunistic Infections in HIV-Infected Adults and Adolescents (available at www.aidsinfo.nih.gov/guidelines). Additional prophylaxis may be indicated based on the cancer treatment and will be indicated as such in the guidelines where appropriate. Measurement of the CD4+ T cell count can be considered more frequently than otherwise required in patients receiving cancer treatments that are anticipated to cause lymphopenia. If febrile neutropenia occurs during cancer treatment, consultation with an infectious disease specialist is strongly recommended.

AIDS-Related Kaposi Sarcoma

Kaposi sarcoma is a multifocal malignancy of endothelial cells, which presents with characteristic red or brown papules. The risk for Kaposi sarcoma in the setting of HIV has been reported to be increased as much as 3640-fold over the general U.S. population,^{6-8,14,50} but this risk

has declined with the ART era.^{6,9,10,15} Still, estimates indicate that the risk of Kaposi sarcoma in PLWH between the years 2009-2012 was elevated approximately 257-fold compared with the general U.S. population,¹⁵ and Kaposi sarcoma accounts for approximately 12% of cancers diagnosed in PLWH.¹¹

Four types of Kaposi sarcoma have been described.^{12,51,52} Classic Kaposi sarcoma generally involves indolent cutaneous lesions, often of the lower extremities, that slowly progress over years to decades. It is most common in older people of Mediterranean, Eastern European, Middle Eastern, and/or Jewish origins. It is much more common in men than in women. Endemic Kaposi sarcoma occurs in children and younger adults (<40 years of age) of equatorial Africa. It is usually more aggressive than classic Kaposi sarcoma, sometimes with visceral, bone, and/or lymph node involvement. When Kaposi sarcoma occurs in the context of immunosuppressive therapy (for organ transplant or other reasons), it is called iatrogenic or transplant-associated Kaposi sarcoma. Although this form of Kaposi sarcoma can be aggressive and involve lymph nodes, mucosa, and/or visceral organs, it frequently responds to a reduction or cessation of immunosuppression. Finally, when Kaposi sarcoma occurs in the setting of HIV seropositivity, it is considered an AIDS-defining illness and is referred to as AIDS-related or epidemic Kaposi sarcoma. When immunosuppression is advanced, AIDS-related Kaposi sarcoma is more common, more aggressive, and more likely to involve viscera and/or lymph nodes than when immunosuppression is minimal. However, AIDS-related Kaposi sarcoma can occur in PLWH with normal CD4+ T cells counts and viral load. Overall, AIDS-related Kaposi sarcoma tends to be more aggressive than other types.

Kaposi sarcoma is highly associated with human herpesvirus 8 infection (HHV-8, also known as Kaposi sarcoma–associated herpesvirus,

KSHV).⁵¹ It is estimated that 95% to 98% of patients with Kaposi sarcoma are seropositive for HHV-8.^{51,52} In a study of 5022 ART-naïve, PLWH enrolled in 6 U.S. randomized clinical trials, 38% were infected with HHV-8.⁵³ HHV-8 infections are usually asymptomatic, and immunosuppression is likely an important factor in the pathogenesis of Kaposi sarcoma in HHV-8–infected individuals. In fact, CD4+ T cell counts and HIV viral load correlate with the risk of Kaposi sarcoma in PLWH.⁵⁴ Thus, effective ART likely lowers the risk of Kaposi sarcoma development. Evidence also suggests that ART improves prognosis of Kaposi sarcoma. The 5-year survival of patients with AIDS-related Kaposi sarcoma has improved in the post-ART era, from 12.1% in 1980-1995 to as high as 88% in the post-ART era.⁵⁵⁻⁵⁷

Diagnosis and Workup of AIDS-Related Kaposi Sarcoma

As described in the guidelines above, AIDS-related Kaposi sarcoma is diagnosed by pathology and immunophenotyping. Workup should include a history and physical exam that includes any history of additional immunosuppression such as transplant or glucocorticoids and HIV testing (if HIV status is unknown). In addition, complete skin and oral exams, with documentation of edema and photography of oral, conjunctival, and cutaneous lesions for documentation of extent of disease are recommended. Referral to an HIV specialist is also recommended, as is care coordination between the HIV specialist and the oncology team (see *HIV Management during Cancer Therapy*, above). All PLWH should have recent T cell subsets including quantitative CD4+ T cell counts and HIV viral load to assess immune function and HIV control. This testing may be done in conjunction with the HIV specialist.

Another essential workup item is chest X-ray to assess for disseminated disease. Depending on symptoms and findings that may be concerning

for possible distant metastases, additional workup may include chest CT with contrast with or without an abdominal/pelvic CT with contrast or an MRI with contrast and/or a PET/CT scan. It is important to note that imaging in PLWH who have cancer is complicated by the increased incidence of non-malignant lesions that may be mistaken for cancer spread or recurrence. Opportunistic infections in the lung include mycobacterium tuberculosis (MTB), cytomegalovirus (CMV), and Pneumocystis jirovecii pneumonia (PCP).⁵⁸ Furthermore, non-infectious, non-malignant pulmonary manifestations of HIV can be difficult to interpret on imaging studies, including interstitial pneumonia and granulomatous disease.^{58,59} Furthermore, brain lesions seen in PLWH may result from opportunistic infections, such as viral encephalitis, aspergillosis, toxoplasmosis, cryptococcosis, bacterial meningitis, tuberculosis, progressive multifocal leukoencephalopathy, and mycobacterium avium complex (MAC).^{60,61} Benign non-infectious brain lesions can also occur in PLWH (eg, vascular complications, hydrocephalus).^{60,61} Similarly, immune response to HIV and opportunistic infections commonly cause lymphadenopathy in PLWH, which can be seen on F-18 FDG PET/CT.^{62,63} Non-malignant causes of lymphadenopathy are more common in patients with higher viral loads and lower CD4+ T cell counts.⁶⁴ Therefore, patients with cancer and HIV infection should have an infectious disease workup for imaging findings, as clinically indicated.

Staging of AIDS-Related Kaposi Sarcoma

As delineated in the guidelines above, AIDS-related Kaposi sarcoma is staged using a TIS system, in which aspects of the tumor (T), immune system (I), and systemic disease (S) are assessed with a 0 for good risk and 1 for poor risk.⁶⁵ However, more recent data has shown that the I stage has less prognostic value than the T or S stages in the presence of ART.⁵⁶ Patients staged as T1S1 appear to have the worst prognosis.

In a study of 211 patients with AIDS-related Kaposi sarcoma, those staged as T1S1 had a 3-year survival rate of 53%, whereas for those staged as T0S0, T1S0, or T0S1, the 3-year survival rates were 88%, 80%, and 81%, respectively ($P = .0001$).⁵⁶

Initial Management of AIDS-Related Kaposi Sarcoma

Patients with limited cutaneous disease that is asymptomatic and cosmetically acceptable to the patient may be treated with ART alone (see below). Those with symptomatic or cosmetically unacceptable limited cutaneous disease should be treated with ART and with the most minimally invasive and least toxic therapy possible. A limited number of cycles of systemic therapy (eg, 3–6; options discussed below) may be sufficient for those initiating or re-initiating ART. Other options include topical treatment, intralesional chemotherapy, radiation, and local excision (all discussed below).

Preferred initial treatment for patients with advanced cutaneous, oral, visceral, or nodal AIDS-related Kaposi sarcoma is ART with clinical trial or systemic therapy. For those not eligible for clinical trial or systemic therapy, radiation can be used with ART. The data supporting these treatment options is described below.

It is important to note that individual Kaposi sarcoma lesions may be distinct clones that arise because of the common risk factors of immunosuppression and persistent HHV-8 infection as opposed to metastases. Furthermore, persistence of HHV-8 infection results in ongoing risk of recurrence/disease progression. Currently eradication of HHV-8 is not possible. Therefore, treatment of existing disease may not prevent occurrence of future lesions, and the goals of therapy are based on disease control.

Antiretroviral Therapy (ART)

Reconstitution of immune function, maintenance of viral suppression, and avoidance of additional immunosuppression are critical to prevent additional Kaposi sarcoma lesions and maintain response to therapy. In fact, in the setting of limited cutaneous disease, remissions or stable disease may occur with optimization of immune function and HIV viral suppression alone.⁶⁶⁻⁷² Therefore, co-management with an HIV specialist to optimize suppression of HIV and reconstitution of immune function with ART is important for patients with AIDS-related Kaposi sarcoma (see *HIV Management during Cancer Therapy*, above).

Initiation of ART may result in immune reconstitution syndrome (IRIS) within 3-6 months in a reported 6% to 39% of patients with AIDS-related Kaposi sarcoma.⁷³⁻⁷⁶ IRIS is characterized by marked lesional swelling, increased tenderness, and peripheral edema. Individuals with pulmonary involvement, the use of glucocorticoids, or advanced immunosuppression may be at increased risk.^{73,74,76} In contrast with management of IRIS for some opportunistic infections, glucocorticoids are contraindicated in Kaposi-sarcoma IRIS because of the potential for life-threatening Kaposi sarcoma exacerbation resulting from stimulatory effects of glucocorticoids on Kaposi sarcoma spindle cells.^{77,78}

Management of Kaposi sarcoma IRIS should involve coordination with an HIV specialist. ART should not be delayed or discontinued unless life-threatening IRIS develops.

Topical Therapies

Topical therapies are an option for patients with limited cutaneous disease that is symptomatic or cosmetically unacceptable. Alitretinoin gel, a retinoid, was studied in a phase III vehicle-controlled, double-blind, multi-centered study, in which 134 patients with AIDS-related Kaposi sarcoma received either 0.1% alitretinoin gel or vehicle gel twice daily for 12 weeks.⁷⁹ The cutaneous tumor response rates were 37% in

the alitretinoin group compared with 7% in the control group. Another very similar randomized, multicenter, double-blind, vehicle-controlled study also compared tumor response rates in patients with AIDS-related Kaposi sarcoma between an alitretinoin group and a control group.⁸⁰ Response rates in the 268 patients were 35% for those receiving 0.1% alitretinoin gel compared with 18% for those who received the vehicle gel. In both of these studies, alitretinoin gel was well tolerated, with mostly mild to moderate adverse events that were limited to the application site and that were relieved when treatment was stopped.

Imiquimod is a topical immune response modulator with antiviral and antitumor activity.⁸¹ It is used in a variety of skin conditions including malignancies and warts.^{81,82} Case reports have shown that imiquimod cream can be safe and effective in some patients with classic or transplant-associated Kaposi sarcoma.⁸³⁻⁸⁸ In a single-center, open-label, phase I/II trial, 17 HIV-negative patients with Kaposi sarcoma received imiquimod 5% cream 3 times per week for 24 weeks.⁸⁹ The response rate was 47%. Over half of the patients reported local itching and erythema, but treatment was generally well tolerated. Imiquimod is not well studied as a treatment for patients with cutaneous AIDS-related Kaposi sarcoma.^{90,91} The panel includes imiquimod as an option for patients with cutaneous AIDS-related Kaposi sarcoma based on extrapolation from the data presented above in other settings, expert opinion, and non-published anecdotal data.

Intralesional Chemotherapy

Intralesional vinblastine is another option for patients with limited mucocutaneous disease that is symptomatic or cosmetically unacceptable. Intralesional injection of vinblastine has been studied in case reports, case series, and one small randomized trial of patients with oral AIDS-related Kaposi sarcoma.⁹²⁻⁹⁸ In a large series of 144 oral Kaposi sarcoma lesions in 50 HIV-positive men, complete response

was seen in 74% of lesions and partial response in 26%.⁹⁵ The recurrence rate was 26%, with a mean disease-free period of 12.9 weeks. Consistent with the safety profile seen in other studies, pain was reported by 72% of participants, ulceration occurred in 22%, and temporary numbness was seen in 12%. Pain is generally mild to moderate and relieved with pain medication, and ulceration is generally self-limiting.

Studies on the use of intralesional vinblastine injection for cutaneous lesions are more limited.^{99,100} In a trial of 11 men with AIDS-related Kaposi sarcoma, 88% of cutaneous lesions showed a complete or partial clinical response.⁹⁹ Treatment resulted in inflammation and blistering of the lesion prior to healing, and the final results were not cosmetically optimal because of post-inflammation hyperpigmentation. Most patients reported aching pain 6 to 48 hours post-treatment that was relieved with pain medication.

Intralesional vinblastine has also been used in cutaneous lesions in patients with classic Kaposi sarcoma.¹⁰¹

Local Excision

Local excision is an option for patients with limited cutaneous disease that is symptomatic or cosmetically unacceptable. However, data regarding outcomes of the excision of cutaneous Kaposi sarcoma lesions are limited and appear to be restricted to HIV-negative individuals.¹⁰²⁻¹⁰⁶

Radiation Therapy

AIDS-related Kaposi sarcoma is radioresponsive, with complete responses rates of treated lesions reported in the range of 68% to 92%.¹⁰⁷⁻¹¹¹ Radiation therapy for AIDS-related Kaposi sarcoma is used in patients with limited cutaneous disease that is symptomatic or

cosmetically unacceptable. For patients with advanced disease, systemic therapy is preferred over radiation therapy in first-line and for relapsed/refractory disease as long as systemic therapy is feasible based on performance status and comorbidities. Radiation in this setting should be reserved for circumstances when systemic therapy is not feasible or when palliative therapy is needed to mitigate pain or other symptoms.¹¹²

When radiation is used, hypofractionated regimens (eg, 20 Gy in 5 fractions) appear to be equally effective to the standard regimen of 24 Gy in 12 fractions.^{113,114} Dose fractionation should be based on the site of treatment with consideration for surrounding normal tissue tolerance.

The side effects of radiation for AIDS-related Kaposi sarcoma are site-dependent, but typically manageable given the low doses needed to achieve a response.¹⁰⁷⁻¹¹⁰ Early recognition and treatment of dermatitis, oral mucositis, and lymphedema are especially important.^{107,109,115} The risk of lymphedema is already elevated in patients with KS and may increase after radiation.¹¹⁶ Early referral to and co-management with a lymphedema specialist is recommended.

Systemic Therapy

The preferred first-line systemic therapy for both limited cutaneous disease and advanced disease is liposomal doxorubicin. In a randomized phase III trial, 258 patients with advanced AIDS-related Kaposi sarcoma were randomized to receive pegylated-liposomal doxorubicin or doxorubicin/bleomycin/vincristine (ABV).¹¹⁷ The overall response rate was 46% (95% CI, 37% to 54%) in the liposomal doxorubicin arm and 25% (95% CI, 17% to 32%) in the ABV arm. The median time to treatment failure was approximately 4 months in both groups. Most patients in both arms experienced ≥1 grade 3/4 adverse event, with leukopenia, nausea/vomiting, anemia, and peripheral

neuropathy as the most common adverse events in the liposomal doxorubicin group. Pegylated-liposomal doxorubicin was also compared with bleomycin/vincristine (BV) in another randomized trial of patients with AIDS-related Kaposi sarcoma (n=241).¹¹⁸ As in the other trial, response rates were superior in the liposomal doxorubicin group compared with the BV group (59% versus 23%; $P < .001$). Pegylated-liposomal doxorubicin resulted in an increased risk of neutropenia, but was less likely to result in early treatment cessation.

Liposomal doxorubicin is associated with risk of cardiotoxicity.¹¹⁹⁻¹²¹ Therefore, a baseline echocardiogram should be performed prior to initial and repeat courses of liposomal doxorubicin, and the lifetime dose should be limited to 400-450 mg/m².

An alternative option for first-line systemic therapy for limited cutaneous and advanced disease is paclitaxel. Early studies showed that it has significant activity in the advanced disease setting, with neutropenia as the most frequent dose-limiting toxicity.^{122,123}

One trial randomized 73 patients with advanced AIDS-related Kaposi sarcoma to paclitaxel or pegylated-liposomal doxorubicin.¹²⁴ The two arms were statistically equivalent with regards to response rates, median progression-free survival, and 2-year survival. A trend toward increase in grade 3 to grade 5 toxicity was seen in the paclitaxel arm (84% vs 66%; $P = .077$), with 1 lethal, grade 5 pulmonary embolism in a patient treated with paclitaxel. A systematic review of randomized trials and observational studies in patients with advanced AIDS-related Kaposi sarcoma found no evident differences between liposomal doxorubicin, liposomal daunorubicin, and paclitaxel, although the number of studies identified was low.¹²⁵

Surveillance of Patients with AIDS-Related Kaposi Sarcoma

Patients treated for AIDS-related Kaposi sarcoma who do not require active treatment and who are without signs of progression should be followed every 3 months for year 1, then every 4-6 months for year 2, then every 6-12 months thereafter. Surveillance should include history and physical (including complete skin and oral exams and documentation of edema and history of additional immunosuppression such as transplant/glucocorticoids), CBC, differential, comprehensive metabolic panel, T cell subsets (CD4+ T-cell count), and HIV viral load. ART compliance should also be assessed. If a change in disease is noted, lesions should be photographed for documentation. Stool testing, chest x-ray or chest CT with contrast, EGD/colonoscopy, and bronchoscopy should be performed only for signs and symptoms concerning for visceral involvement or, in the case of progression/refractory disease, before a new therapy is initiated.

It is important to note that HHV-8 is not eradicated with treatment of Kaposi sarcoma, and the risk of future Kaposi sarcoma persists even after complete remission. Optimization and monitoring of HIV control and immune function is important to minimize this risk, because disease risk generally decreases with immune reconstitution. However, Kaposi sarcoma can persist, relapse, or present even in the setting of normal values of T cell subsets. Less frequent (every 6-12 months) oncologic monitoring may be appropriate for select patients with undetectable HIV viral loads, normal T cell subsets, and Kaposi sarcoma that is stable for ≥ 2 years, providing the patient has regular follow up with an HIV specialist.

Systemic Therapy of Relapsed/Refractory Disease

At first progression, the same systemic therapy options as in first line (liposomal doxorubicin and paclitaxel, discussed above) may be considered as follows:

- If first-line therapy was tolerated and a durable response (>3 months) was seen, then a repeat of the therapy used in first line can be considered.
- If there was no response to first-line systemic therapy, then an alternative first-line therapy option should be given.

Following subsequent progressions, liposomal doxorubicin or paclitaxel, whichever has not yet been administered, is recommended.^{126,127} In third-line, the panel recommends pomalidomide as the preferred regimen. Pomalidomide was studied in a phase I/II trial of 7 HIV-negative and 15 PLWH with Kaposi sarcoma.¹²⁸ PLWH were required to have viremia controlled and either progressive or stable Kaposi sarcoma on ART. Most of the participants (19 of 22) had previous therapy for Kaposi sarcoma, exclusive of ART. The response rate was 60% in the HIV-infected group (95% CI, 32%-84%). Grade 3/4 adverse events that might have occurred due to pomalidomide were neutropenia, infection, and edema.

Other treatment options for subsequent lines of therapy for relapsed/refractory disease, listed in alphabetical order, include bevacizumab, etoposide, gemcitabine, imatinib, interferon, nab-paclitaxel, thalidomide, and vinorelbine, but data for these agents are generally limited, as described below.

Bevacizumab was assessed in a phase II study of 17 PLWH with Kaposi sarcoma who had progressive or stable disease on ART.¹²⁹

Thirteen of the patients had received prior chemotherapy for Kaposi sarcoma. The complete response rate was 19% and the partial

response rate was 12%, for an overall response rate of 31% (95% CI, 11% to 59%). Adverse events included hypertension (n=7), neutropenia (n=5), cellulitis (n=3), and headache (n=2).

Etoposide has been studied in multiple phase II trials of patients with AIDS-related Kaposi sarcoma.¹³⁰⁻¹³² In one of these trials, 36 patients with previously treated AIDS-related Kaposi sarcoma received a course of oral etoposide, and the overall response rate was 36%, with stable disease occurring in 33% of the participants.¹³² The median duration of response was about 6 months. Grade 3/4 neutropenia occurred in 28%, and opportunistic infections in 22%. The other trials also showed oral etoposide to have clinical activity and be fairly well tolerated.

Evidence for the use of gemcitabine in patients with refractory AIDS-related Kaposi sarcoma comes only from a retrospective analysis of 23 patients who had been treated with first-line ABV.¹³³ Complete response was seen in 3 patients (13%), partial response in 8 (35%), and stable disease in 11 (48%). Only 1 patient had progressive disease. Grade 3/4 adverse events include leukopenia, pain, fatigue, and neutropenia.

Imatinib appears to have activity in AIDS-related Kaposi sarcoma.^{134,135} The strongest evidence comes from a multicenter phase II trial, in which 30 patients were treated with imatinib.¹³⁶ 18 patients (60%) had received prior therapy. Although no complete responses were seen, 33% achieved partial response and 20% had stable disease. The median duration of response was approximately 8 months, with disease progression in 7 patients (23%). Grade 3/4 adverse events attributed to imatinib included allergic reaction/hypersensitivity, nausea, dehydration, and cellulitis, but only five patients (17%) discontinued therapy because of adverse events.

Early studies suggested that various forms of interferon had clinical activity in AIDS-related Kaposi sarcoma.¹³⁷⁻¹⁴⁰ Several studies in the post-ART era have focused specifically on interferon alpha-2b in this population.^{141,142} In one randomized phase II trial, the safety and efficacy of low-dose interferon alpha-2b was assessed in 35 patients with AIDS-related cutaneous Kaposi sarcoma.¹⁴¹ The response rate was 40%, and the median duration of response was approximately 25 months. Grade 3/4 neutropenia occurred in 3% of patients.

Evidence for the use of nab-paclitaxel in Kaposi sarcoma appears to be limited to 1 abstract of a phase II trial of 6 patients with classic Kaposi sarcoma.¹⁴³ Partial (n=2) or complete responses (n=4) were seen in all patients. Grade 3 adverse events were neutropenia in half of the patients and thrombocytopenia in 1 of the 6.

Thalidomide has been studied in AIDS-related Kaposi sarcoma in 2 phase II trials.^{144,145} One of these trials included 17 assessable patients with progressive disease.¹⁴⁴ Partial responses were seen in 47%, and stable disease was seen in 12%. Time to progression was a median 7.3 months. The most frequently reported side effects were drowsiness in 45% of participants and depression in 35%.

Evidence for the activity of vinorelbine in AIDS-related Kaposi sarcoma comes from a phase II trial of 35 assessable patients with progressive disease.¹⁴⁶ Complete clinical responses were seen in 9%, and partial responses were seen in 34%. The median duration of response was about 6 months. Neutropenia was the most frequent dose-limiting toxicity, but other side effects were mild and reversible and the treatment was generally well tolerated.

Summary

Management of AIDS-related Kaposi sarcoma depends on location and extent of disease. Patients with limited cutaneous disease that is asymptomatic and cosmetically acceptable to the patient may be treated with ART alone. Remissions or stable disease may occur with optimization of immune function and HIV viral suppression alone.

Those with symptomatic or cosmetically unacceptable limited cutaneous disease should be treated with ART and with the most minimally invasive and least toxic therapy possible. Options include a limited number of cycles of systemic therapy, topical treatment, intralesional chemotherapy, radiation, and local excision.

Preferred initial treatment for patients with advanced cutaneous, oral, visceral, or nodal AIDS-related Kaposi sarcoma is ART with clinical trial or systemic therapy. For those not eligible for clinical trial or systemic therapy, radiation can be used with ART. As lymphedema often complicates Kaposi sarcoma, early involvement of a lymphedema specialist is recommended.

Surveillance of patients treated for AIDS-related Kaposi sarcoma is important, as disease can recur after an initial complete response and in the setting of normal values of T-cell subsets. Persistence of HHV-8 and emergence of distinct tumor clones can lead to disease progression and relapse. Furthermore, because individual Kaposi sarcoma lesions are often distinct clones as opposed to metastases, treatment of existing disease does not prevent occurrence of new lesions.

For relapsed/refractory disease, a typical systemic therapy sequence would be first-line liposomal doxorubicin, followed by second-line paclitaxel, followed by pomalidomide in the third-line of treatment.



NCCN Guidelines Version 1.2018 AIDS-Related Kaposi Sarcoma

Additional lines of other therapies can be given, and the any systemic therapy that was tolerated with a durable response can be repeated.

Overall, the survival of patients with AIDS-related Kaposi sarcoma has improved greatly, and long-term survival can be the goal for many patients. However, the goals of therapy for patients with advanced disease are namely reducing or reversing symptoms and mitigating end organ damage. Complete remissions in this setting are rare, but effective therapy can lead to long-term disease control.

References

1. U.S. Statistics. HIV.gov; 2017. Available at: <https://www.hiv.gov/hiv-basics/overview/data-and-trends/statistics>. Accessed August 16, 2017.
2. Goedert JJ, Cote TR, Virgo P, et al. Spectrum of AIDS-associated malignant disorders. *Lancet* 1998;351:1833-1839. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9652666>.
3. Frisch M, Biggar RJ, Engels EA, et al. Association of cancer with AIDS-related immunosuppression in adults. *JAMA* 2001;285:1736-1745. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11277828>.
4. Detels R, Munoz A, McFarlane G, et al. Effectiveness of potent antiretroviral therapy on time to AIDS and death in men with known HIV infection duration. Multicenter AIDS Cohort Study Investigators. *JAMA* 1998;280:1497-1503. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9809730>.
5. Palella FJ, Jr., Delaney KM, Moorman AC, et al. Declining morbidity and mortality among patients with advanced human immunodeficiency virus infection. HIV Outpatient Study Investigators. *N Engl J Med* 1998;338:853-860. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9516219>.
6. Engels EA, Pfeiffer RM, Goedert JJ, et al. Trends in cancer risk among people with AIDS in the United States 1980-2002. *AIDS* 2006;20:1645-1654. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16868446>.
7. Engels EA, Biggar RJ, Hall HI, et al. Cancer risk in people infected with human immunodeficiency virus in the United States. *Int J Cancer* 2008;123:187-194. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18435450>.
8. Patel P, Hanson DL, Sullivan PS, et al. Incidence of types of cancer among HIV-infected persons compared with the general population in the United States, 1992-2003. *Ann Intern Med* 2008;148:728-736. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18490686>.
9. Robbins HA, Shiels MS, Pfeiffer RM, Engels EA. Epidemiologic contributions to recent cancer trends among HIV-infected people in the United States. *AIDS* 2014;28:881-890. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24300545>.
10. Shiels MS, Pfeiffer RM, Hall HI, et al. Proportions of Kaposi sarcoma, selected non-Hodgkin lymphomas, and cervical cancer in the United States occurring in persons with AIDS, 1980-2007. *JAMA* 2011;305:1450-1459. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21486978>.
11. Robbins HA, Pfeiffer RM, Shiels MS, et al. Excess cancers among HIV-infected people in the United States. *J Natl Cancer Inst* 2015;107. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25663691>.
12. Angeletti PC, Zhang L, Wood C. The viral etiology of AIDS-associated malignancies. *Adv Pharmacol* 2008;56:509-557. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18086422>.
13. Chaturvedi AK, Madeleine MM, Biggar RJ, Engels EA. Risk of human papillomavirus-associated cancers among persons with AIDS. *J Natl Cancer Inst* 2009;101:1120-1130. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19648510>.
14. Grulich AE, van Leeuwen MT, Falster MO, Vajdic CM. Incidence of cancers in people with HIV/AIDS compared with immunosuppressed transplant recipients: a meta-analysis. *Lancet* 2007;370:59-67. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17617273>.
15. Hernandez-Ramirez RU, Shiels MS, Dubrow R, Engels EA. Cancer risk in HIV-infected people in the USA from 1996 to 2012: a population-based, registry-linkage study. *Lancet HIV* 2017. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28803888>.
16. Meijide H, Pertega S, Rodriguez-Ororio I, et al. Increased incidence of cancer observed in HIV/HCV-coinfected patients versus HIV-monoinfected, 1993-2014. *AIDS* 2017. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28328796>.

17. Tesoriero JM, Gieryic SM, Carrascal A, Lavigne HE. Smoking among HIV positive New Yorkers: prevalence, frequency, and opportunities for cessation. *AIDS Behav* 2010;14:824-835. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18777131>.
18. Helleberg M, Afzal S, Kronborg G, et al. Mortality attributable to smoking among HIV-1-infected individuals: a nationwide, population-based cohort study. *Clin Infect Dis* 2013;56:727-734. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23254417>.
19. McGinnis KA, Fultz SL, Skanderson M, et al. Hepatocellular carcinoma and non-Hodgkin's lymphoma: the roles of HIV, hepatitis C infection, and alcohol abuse. *J Clin Oncol* 2006;24:5005-5009. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17075119>.
20. Park LS, Hernandez-Ramirez RU, Silverberg MJ, et al. Prevalence of non-HIV cancer risk factors in persons living with HIV/AIDS: a meta-analysis. *AIDS* 2016;30:273-291. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26691548>.
21. Rentsch C, Tate JP, Akgun KM, et al. Alcohol-related diagnoses and all-cause hospitalization among HIV-infected and uninfected patients: a longitudinal analysis of United States veterans from 1997 to 2011. *AIDS Behav* 2016;20:555-564. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25711299>.
22. Marks G, Crepaz N, Senterfitt JW, Janssen RS. Meta-analysis of high-risk sexual behavior in persons aware and unaware they are infected with HIV in the United States: implications for HIV prevention programs. *J Acquir Immune Defic Syndr* 2005;39:446-453. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16010168>.
23. Branson BM, Handsfield HH, Lampe MA, et al. Revised recommendations for HIV testing of adults, adolescents, and pregnant women in health-care settings. *MMWR Recomm Rep* 2006;55:1-17; quiz CE11-14. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16988643>.
24. Chiao EY, Dezube BJ, Krown SE, et al. Time for oncologists to opt in for routine opt-out HIV testing? *JAMA* 2010;304:334-339. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20639567>.
25. Hwang JP, Granwehr BP, Torres HA, et al. HIV testing in patients with cancer at the initiation of therapy at a large US comprehensive cancer center. *J Oncol Pract* 2015;11:384-390. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26243649>.
26. Li J, Thompson TD, Tai E, et al. Testing for human immunodeficiency virus among cancer survivors under age 65 in the United States. *Prev Chronic Dis* 2014;11:E200. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25393748>.
27. HIV Care Continuum. HIV.gov; 2017. Available at: <https://www.hiv.gov/federal-response/policies-issues/hiv-aids-care-continuum>. Accessed August 16, 2017.
28. Flash CA, Pasalar S, Hemmige V, et al. Benefits of a routine opt-out HIV testing and linkage to care program for previously diagnosed patients in publicly funded emergency departments in Houston, TX. *J Acquir Immune Defic Syndr* 2015;69 Suppl 1:S8-15. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25867782>.
29. Irvine MK, Chamberlin SA, Robbins RS, et al. Improvements in HIV care engagement and viral load suppression following enrollment in a comprehensive HIV care coordination program. *Clin Infect Dis* 2015;60:298-310. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25301208>.
30. Kitahata MM, Gange SJ, Abraham AG, et al. Effect of early versus deferred antiretroviral therapy for HIV on survival. *N Engl J Med* 2009;360:1815-1826. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19339714>.
31. El-Sadr WM, Lundgren J, Neaton JD, et al. CD4+ count-guided interruption of antiretroviral treatment. *N Engl J Med* 2006;355:2283-2296. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17135583>.

32. Hessel NA, Pipkin S, Schwarcz S, et al. The impact of highly active antiretroviral therapy on non-AIDS-defining cancers among adults with AIDS. *Am J Epidemiol* 2007;165:1143-1153. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17344204>.

33. Gerard L, Galicier L, Maillard A, et al. Systemic non-Hodgkin lymphoma in HIV-infected patients with effective suppression of HIV replication: persistent occurrence but improved survival. *J Acquir Immune Defic Syndr* 2002;30:478-484. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12154338>.

34. Torres HA, Mulanovich V. Management of HIV infection in patients with cancer receiving chemotherapy. *Clin Infect Dis* 2014;59:106-114. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24642555>.

35. Buchacz K, Baker RK, Palella FJ, Jr., et al. AIDS-defining opportunistic illnesses in US patients, 1994-2007: a cohort study. *AIDS* 2010;24:1549-1559. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20502317>.

36. Ledergerber B, Egger M, Erard V, et al. AIDS-related opportunistic illnesses occurring after initiation of potent antiretroviral therapy: the Swiss HIV Cohort Study. *JAMA* 1999;282:2220-2226. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10605973>.

37. Schwarcz L, Chen MJ, Vittinghoff E, et al. Declining incidence of AIDS-defining opportunistic illnesses: results from 16 years of population-based AIDS surveillance. *AIDS* 2013;27:597-605. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23079812>.

38. Djawe K, Buchacz K, Hsu L, et al. Mortality risk after AIDS-defining opportunistic illness among HIV-infected persons--San Francisco, 1981-2012. *J Infect Dis* 2015;212:1366-1375. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26044289>.

39. Borg C, Ray-Coquard I, Philip I, et al. CD4 lymphopenia as a risk factor for febrile neutropenia and early death after cytotoxic

chemotherapy in adult patients with cancer. *Cancer* 2004;101:2675-2680. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15503313>.

40. Dale DC, McCarter GC, Crawford J, Lyman GH. Myelotoxicity and dose intensity of chemotherapy: reporting practices from randomized clinical trials. *J Natl Compr Canc Netw* 2003;1:440-454. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19761076>.

41. Seropian S, Nadkarni R, Jillella AP, et al. Neutropenic infections in 100 patients with non-Hodgkin's lymphoma or Hodgkin's disease treated with high-dose BEAM chemotherapy and peripheral blood progenitor cell transplant: out-patient treatment is a viable option. *Bone Marrow Transplant* 1999;23:599-605. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10217191>.

42. Savage DG, Lindenbaum J, Garrett TJ. Biphasic pattern of bacterial infection in multiple myeloma. *Ann Intern Med* 1982;96:47-50. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/6976144>.

43. Griffiths H, Lea J, Bunch C, et al. Predictors of infection in chronic lymphocytic leukaemia (CLL). *Clin Exp Immunol* 1992;89:374-377. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/1516254>.

44. Bodey GP, Buckley M, Sathe YS, Freireich EJ. Quantitative relationships between circulating leukocytes and infection in patients with acute leukemia. *Ann Intern Med* 1966;64:328-340. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/5216294>.

45. Morrison VA. Immunosuppression associated with novel chemotherapy agents and monoclonal antibodies. *Clin Infect Dis* 2014;59 Suppl 5:S360-364. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25352632>.

46. Schimpff SC. Empiric antibiotic therapy for granulocytopenic cancer patients. *Am J Med* 1986;80:13-20. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/3521270>.

47. Park J, Kim TM, Hwang JH, et al. Risk factors for febrile neutropenia during chemotherapy for HIV-related lymphoma. *J Korean Med Sci* 2012;27:1468-1471. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23255844>.
48. Sparano JA, Hu X, Wiernik PH, et al. Opportunistic infection and immunologic function in patients with human immunodeficiency virus-associated non-Hodgkin's lymphoma treated with chemotherapy. *J Natl Cancer Inst* 1997;89:301-307. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9048834>.
49. Wang ES, Straus DJ, Teruya-Feldstein J, et al. Intensive chemotherapy with cyclophosphamide, doxorubicin, high-dose methotrexate/ifosfamide, etoposide, and high-dose cytarabine (CODOX-M/IVAC) for human immunodeficiency virus-associated Burkitt lymphoma. *Cancer* 2003;98:1196-1205. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12973843>.
50. Lee JY, Dhakal I, Casper C, et al. Risk of cancer among commercially insured HIV-infected adults on antiretroviral therapy. *J Cancer Epidemiol* 2016;2016:2138259. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27882054>.
51. Antman K, Chang Y. Kaposi's sarcoma. *N Engl J Med* 2000;342:1027-1038. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10749966>.
52. Hiatt KM, Nelson AM, Lichy JH, Fanburg-Smith JC. Classic Kaposi Sarcoma in the United States over the last two decades: a clinicopathologic and molecular study of 438 non-HIV-related Kaposi Sarcoma patients with comparison to HIV-related Kaposi Sarcoma. *Mod Pathol* 2008;21:572-582. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18376387>.
53. Labo N, Miley W, Benson CA, et al. Epidemiology of Kaposi's sarcoma-associated herpesvirus in HIV-1-infected US persons in the era of combination antiretroviral therapy. *AIDS* 2015;29:1217-1225. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26035321>.
54. Dubrow R, Qin L, Lin H, et al. Association of CD4+ T-cell count, HIV-1 RNA viral load, and antiretroviral therapy with Kaposi sarcoma risk among HIV-infected persons in the United States and Canada. *J Acquir Immune Defic Syndr* 2017;75:382-390. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28394855>.
55. Armstrong AW, Lam KH, Chase EP. Epidemiology of classic and AIDS-related Kaposi's sarcoma in the USA: incidence, survival, and geographical distribution from 1975 to 2005. *Epidemiol Infect* 2013;141:200-206. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22404880>.
56. Nasti G, Talamini R, Antinori A, et al. AIDS-related Kaposi's Sarcoma: evaluation of potential new prognostic factors and assessment of the AIDS Clinical Trial Group Staging System in the Haart Era--the Italian Cooperative Group on AIDS and Tumors and the Italian Cohort of Patients Naive From Antiretrovirals. *J Clin Oncol* 2003;21:2876-2882. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12885804>.
57. Royse KE, El Chaer F, Amirian ES, et al. Disparities in Kaposi sarcoma incidence and survival in the United States: 2000-2013. *PLoS One* 2017;12:e0182750. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28829790>.
58. Allen CM, Al-Jahdali HH, Irion KL, et al. Imaging lung manifestations of HIV/AIDS. *Ann Thorac Med* 2010;5:201-216. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20981180>.
59. Gingo MR, Morris A. Pathogenesis of HIV and the lung. *Curr HIV/AIDS Rep* 2013;10:42-50. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23079728>.
60. Gottumukkala RV, Romero JM, Riascos RF, et al. Imaging of the brain in patients with human immunodeficiency virus infection. *Top Magn Reson Imaging* 2014;23:275-291. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25296273>.



NCCN Guidelines Version 1.2018

AIDS-Related Kaposi Sarcoma

61. Langford TD, Letendre SL, Larrea GJ, Masliah E. Changing patterns in the neuropathogenesis of HIV during the HAART era. *Brain Pathol* 2003;13:195-210. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12744473>.
62. Brust D, Polis M, Davey R, et al. Fluorodeoxyglucose imaging in healthy subjects with HIV infection: impact of disease stage and therapy on pattern of nodal activation. *AIDS* 2006;20:985-993. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16603850>.
63. Scharko AM, Perlman SB, Pyzalski RW, et al. Whole-body positron emission tomography in patients with HIV-1 infection. *Lancet* 2003;362:959-961. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/14511930>.
64. Goshen E, Davidson T, Avigdor A, et al. PET/CT in the evaluation of lymphoma in patients with HIV-1 with suppressed viral loads. *Clin Nucl Med* 2008;33:610-614. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18716509>.
65. Krown SE, Metroka C, Wernz JC. Kaposi's sarcoma in the acquired immune deficiency syndrome: a proposal for uniform evaluation, response, and staging criteria. AIDS Clinical Trials Group Oncology Committee. *J Clin Oncol* 1989;7:1201-1207. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/2671281>.
66. Beatrous SV, Grisoli SB, Riahi RR, Cohen PR. Cutaneous HIV-associated Kaposi sarcoma: a potential setting for management by clinical observation. *Dermatol Online J* 2017;23. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28633734>.
67. Asiimwe F, Moore D, Were W, et al. Clinical outcomes of HIV-infected patients with Kaposi's sarcoma receiving nonnucleoside reverse transcriptase inhibitor-based antiretroviral therapy in Uganda. *HIV Med* 2012;13:166-171. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22112164>.
68. Bower M, Weir J, Francis N, et al. The effect of HAART in 254 consecutive patients with AIDS-related Kaposi's sarcoma. *AIDS* 2009;23:1701-1706. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19550283>.
69. Cattelan AM, Calabro ML, Gasperini P, et al. Acquired immunodeficiency syndrome-related Kaposi's sarcoma regression after highly active antiretroviral therapy: biologic correlates of clinical outcome. *J Natl Cancer Inst Monogr* 2001:44-49. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11158206>.
70. Dupont C, Vasseur E, Beauchet A, et al. Long-term efficacy on Kaposi's sarcoma of highly active antiretroviral therapy in a cohort of HIV-positive patients. CISH 92. Centre d'information et de soins de l'immunodeficiency humaine. *AIDS* 2000;14:987-993. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10853980>.
71. Nguyen HQ, Magaret AS, Kitahata MM, et al. Persistent Kaposi sarcoma in the era of highly active antiretroviral therapy: characterizing the predictors of clinical response. *AIDS* 2008;22:937-945. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18453853>.
72. Mosam A, Shaik F, Uldrick TS, et al. A randomized controlled trial of highly active antiretroviral therapy versus highly active antiretroviral therapy and chemotherapy in therapy-naive patients with HIV-associated Kaposi sarcoma in South Africa. *J Acquir Immune Defic Syndr* 2012;60:150-157. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22395672>.
73. Bower M, Nelson M, Young AM, et al. Immune reconstitution inflammatory syndrome associated with Kaposi's sarcoma. *J Clin Oncol* 2005;23:5224-5228. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16051964>.
74. Fernandez-Sanchez M, Iglesias MC, Ablanedo-Terrazas Y, et al. Steroids are a risk factor for Kaposi's sarcoma-immune reconstitution inflammatory syndrome and mortality in HIV infection. *AIDS*



2016;30:909-914. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/26636923>.

75. Letang E, Almeida JM, Miro JM, et al. Predictors of immune reconstitution inflammatory syndrome-associated with kaposi sarcoma in mozambique: a prospective study. *J Acquir Immune Defic Syndr* 2010;53:589-597. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/19801945>.

76. Volkow P, Cesarman-Maus G, Garciadiego-Fossas P, et al. Clinical characteristics, predictors of immune reconstitution inflammatory syndrome and long-term prognosis in patients with Kaposi sarcoma. *AIDS Res Ther* 2017;14:30. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28558783>.

77. Guo WX, Antakly T. AIDS-related Kaposi's sarcoma: evidence for direct stimulatory effect of glucocorticoid on cell proliferation. *Am J Pathol* 1995;146:727-734. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/7887453>.

78. Volkow PF, Cornejo P, Zinser JW, et al. Life-threatening exacerbation of Kaposi's sarcoma after prednisone treatment for immune reconstitution inflammatory syndrome. *AIDS* 2008;22:663-665. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18317012>.

79. Bodsworth NJ, Bloch M, Bower M, et al. Phase III vehicle-controlled, multi-centered study of topical alitretinoin gel 0.1% in cutaneous AIDS-related Kaposi's sarcoma. *Am J Clin Dermatol* 2001;2:77-87. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11705307>.

80. Walmsley S, Northfelt DW, Melosky B, et al. Treatment of AIDS-related cutaneous Kaposi's sarcoma with topical alitretinoin (9-cis-retinoic acid) gel. Panretin Gel North American Study Group. *J Acquir Immune Defic Syndr* 1999;22:235-246. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/10770343>.

81. Bubna AK. Imiquimod - Its role in the treatment of cutaneous malignancies. *Indian J Pharmacol* 2015;47:354-359. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26288465>.

82. Ganjian S, Ourian AJ, Shamtoub G, et al. Off-label indications for imiquimod. *Dermatol Online J* 2009;15:4. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19624982>.

83. Babel N, Eibl N, Ulrich C, et al. Development of Kaposi's sarcoma under sirolimus-based immunosuppression and successful treatment with imiquimod. *Transpl Infect Dis* 2008;10:59-62. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17428275>.

84. Benomar S, Boutayeb S, Benzekri L, et al. Kaposi's sarcoma responding to topical imiquimod 5% cream: a case report. *Cases J* 2009;2:7092. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20181188>.

85. Bernardini B, Faggion D, Calabro L, et al. Imiquimod for the treatment of classical Kaposi's sarcoma. *Acta Derm Venereol* 2010;90:417-418. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20574613>.

86. Prinz Vavricka BM, Hofbauer GF, Dummer R, et al. Topical treatment of cutaneous Kaposi sarcoma with imiquimod 5% in renal-transplant recipients: a clinicopathological observation. *Clin Exp Dermatol* 2012;37:620-625. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22300351>.

87. Gunduz K, Gunay U, Inanir I, et al. Efficacy of 5% imiquimod cream in a patient with classic Kaposi sarcoma. *J Dermatol Case Rep* 2012;6:52-53. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22826720>.

88. Fairley JL, Denham I, Yoganathan S, Read TR. Topical imiquimod 5% as a treatment for localized genital Kaposi's sarcoma in an HIV-negative man: a case report. *Int J STD AIDS* 2012;23:907-908. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23258836>.



89. Celestin Schartz NE, Chevret S, Paz C, et al. Imiquimod 5% cream for treatment of HIV-negative Kaposi's sarcoma skin lesions: A phase I to II, open-label trial in 17 patients. *J Am Acad Dermatol* 2008;58:585-591. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18068265>.

90. Rosen T. Limited extent AIDS-related cutaneous Kaposi's sarcoma responsive to imiquimod 5% cream. *Int J Dermatol* 2006;45:854-856. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16863526>.

91. Lebari D, Gohil J, Patnaik L, Wasef W. Isolated penile Kaposi's sarcoma in a HIV-positive patient stable on treatment for three years. *Int J STD AIDS* 2014;25:607-610. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24492851>.

92. Epstein JB, Lozada-Nur F, McLeod WA, Spinelli J. Oral Kaposi's sarcoma in acquired immunodeficiency syndrome. Review of management and report of the efficacy of intralesional vinblastine. *Cancer* 1989;64:2424-2430. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/2819653>.

93. Epstein JB. Treatment of oral Kaposi sarcoma with intralesional vinblastine. *Cancer* 1993;71:1722-1725. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8448735>.

94. Ramirez-Amador V, Esquivel-Pedraza L, Lozada-Nur F, et al. Intralesional vinblastine vs. 3% sodium tetradecyl sulfate for the treatment of oral Kaposi's sarcoma. A double blind, randomized clinical trial. *Oral Oncol* 2002;38:460-467. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12110340>.

95. Flaitz CM, Nichols CM, Hicks MJ. Role of intralesional vinblastine administration in treatment of intraoral Kaposi's sarcoma in AIDS. *Eur J Cancer B Oral Oncol* 1995;31B:280-285. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/7492927>.

96. Friedman M, Venkatesan TK, Caldarelli DD. Intralesional vinblastine for treating AIDS-associated Kaposi's sarcoma of the oropharynx and

larynx. *Ann Otol Rhinol Laryngol* 1996;105:272-274. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8604887>.

97. McCormick SU. Intralesional vinblastine injections for the treatment of oral Kaposi's sarcoma: report of 10 patients with 2-year follow-up. *J Oral Maxillofac Surg* 1996;54:583-587; discussion 588-589. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8632242>.

98. Shimomura S, Kikuchi Y, Oka S, Ishitoya J. Local treatment of AIDS-associated bulky Kaposi's sarcoma in the head and neck region. *Auris Nasus Larynx* 2000;27:335-338. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10996492>.

99. Boudreaux AA, Smith LL, Cosby CD, et al. Intralesional vinblastine for cutaneous Kaposi's sarcoma associated with acquired immunodeficiency syndrome. A clinical trial to evaluate efficacy and discomfort associated with infection. *J Am Acad Dermatol* 1993;28:61-65. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8381146>.

100. Smith KJ, Skelton HG, Turiansky G, Wagner KF. Hyaluronidase enhances the therapeutic effect of vinblastine in intralesional treatment of Kaposi's sarcoma. Military Medical Consortium for the Advancement of Retroviral Research (MMCARR). *J Am Acad Dermatol* 1997;36:239-242. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9039176>.

101. Vassallo C, Carugno A, Derlino F, et al. Intralesional vinblastine injections for treatment of classic Kaposi sarcoma in diabetic patients. *Cutis* 2015;95:E28-34. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26057517>.

102. Sen F, Tambas M, Ciftci R, et al. Factors affecting progression-free survival in non-HIV-related Kaposi sarcoma. *J Dermatolog Treat* 2016;27:275-277. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26368051>.

103. Schmidt BM, Holmes CM. Classic solitary Kaposi sarcoma of the foot in an immunocompetent patient: a case report. *Wounds*



NCCN Guidelines Version 1.2018

AIDS-Related Kaposi Sarcoma

2016;28:E35-E40. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27701130>.

104. Cecchi R, Troiano M, Ghilardi M, Bartoli L. Kaposi sarcoma of the penis in an HIV-negative patient. *J Cutan Med Surg* 2011;15:118-120.

Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21477561>.

105. Ozbek MR, Kutlu N. A rare case of Kaposi's sarcoma; hand localization. *Handchir Mikrochir Plast Chir* 1990;22:107-109. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/2338301>.

106. Weintraub CM, Skudowitz RB. Excision of 1,674 classic Kaposi's sarcomas. *S Afr J Surg* 2002;40:80. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/12162235>.

107. Becker G, Bottke D. Radiotherapy in the management of Kaposi's sarcoma. *Onkologie* 2006;29:329-333. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/16874018>.

108. Caccialanza M, Marca S, Piccinno R, Eulisse G. Radiotherapy of classic and human immunodeficiency virus-related Kaposi's sarcoma: results in 1482 lesions. *J Eur Acad Dermatol Venereol* 2008;22:297-302. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18269597>.

109. Cooper JS, Steinfeld AD, Lerch I. Intentions and outcomes in the radiotherapeutic management of epidemic Kaposi's sarcoma. *Int J Radiat Oncol Biol Phys* 1991;20:419-422. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/1995526>.

110. Donato V, Guarnaccia R, Dognini J, et al. Radiation therapy in the treatment of HIV-related Kaposi's sarcoma. *Anticancer Res* 2013;33:2153-2157. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23645769>.

111. Kirova YM, Belembaogo E, Frikha H, et al. Radiotherapy in the management of epidemic Kaposi's sarcoma: a retrospective study of 643 cases. *Radiother Oncol* 1998;46:19-22. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/9488122>.

112. Nobler MP, Leddy ME, Huh SH. The impact of palliative irradiation on the management of patients with acquired immune deficiency syndrome. *J Clin Oncol* 1987;5:107-112. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/2433407>.

113. Singh NB, Lakier RH, Donde B. Hypofractionated radiation therapy in the treatment of epidemic Kaposi sarcoma--a prospective randomized trial. *Radiother Oncol* 2008;88:211-216. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/18439694>.

114. Tsao MN, Sinclair E, Assaad D, et al. Radiation therapy for the treatment of skin Kaposi sarcoma. *Ann Palliat Med* 2016;5:298-302. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27701876>.

115. Wang J, Boerma M, Fu Q, Hauer-Jensen M. Radiation responses in skin and connective tissues: effect on wound healing and surgical outcome. *Hernia* 2006;10:502-506. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/17047884>.

116. Spalek M. Chronic radiation-induced dermatitis: challenges and solutions. *Clin Cosmet Investig Dermatol* 2016;9:473-482. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28003769>.

117. Northfelt DW, Dezube BJ, Thommes JA, et al. Pegylated-liposomal doxorubicin versus doxorubicin, bleomycin, and vincristine in the treatment of AIDS-related Kaposi's sarcoma: results of a randomized phase III clinical trial. *J Clin Oncol* 1998;16:2445-2451. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/9667262>.

118. Stewart S, Jablonowski H, Goebel FD, et al. Randomized comparative trial of pegylated liposomal doxorubicin versus bleomycin and vincristine in the treatment of AIDS-related Kaposi's sarcoma. International Pegylated Liposomal Doxorubicin Study Group. *J Clin Oncol* 1998;16:683-691. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/9469358>.

119. Armenian SH, Lacchetti C, Barac A, et al. Prevention and monitoring of cardiac dysfunction in survivors of adult cancers:

American Society of Clinical Oncology clinical practice guideline. J Clin Oncol 2017;35:893-911. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27918725>.

120. Smith LA, Cornelius VR, Plummer CJ, et al. Cardiotoxicity of anthracycline agents for the treatment of cancer: systematic review and meta-analysis of randomised controlled trials. BMC Cancer 2010;10:337. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20587042>.

121. Carver JR, Shapiro CL, Ng A, et al. American Society of Clinical Oncology clinical evidence review on the ongoing care of adult cancer survivors: cardiac and pulmonary late effects. J Clin Oncol 2007;25:3991-4008. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17577017>.

122. Welles L, Saville MW, Lietzau J, et al. Phase II trial with dose titration of paclitaxel for the therapy of human immunodeficiency virus-associated Kaposi's sarcoma. J Clin Oncol 1998;16:1112-1121. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9508198>.

123. Saville MW, Lietzau J, Pluda JM, et al. Treatment of HIV-associated Kaposi's sarcoma with paclitaxel. Lancet 1995;346:26-28. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/7603142>.

124. Cianfrocca M, Lee S, Von Roenn J, et al. Randomized trial of paclitaxel versus pegylated liposomal doxorubicin for advanced human immunodeficiency virus-associated Kaposi sarcoma: evidence of symptom palliation from chemotherapy. Cancer 2010;116:3969-3977. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20564162>.

125. Gbabe OF, Okwundu CI, Dedicoat M, Freeman EE. Treatment of severe or progressive Kaposi's sarcoma in HIV-infected adults. Cochrane Database Syst Rev 2014:CD003256. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25313415>.

126. Northfelt DW, Dezube BJ, Thommes JA, et al. Efficacy of pegylated-liposomal doxorubicin in the treatment of AIDS-related

Kaposi's sarcoma after failure of standard chemotherapy. J Clin Oncol 1997;15:653-659. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9053490>.

127. Stebbing J, Wildfire A, Portsmouth S, et al. Paclitaxel for anthracycline-resistant AIDS-related Kaposi's sarcoma: clinical and angiogenic correlations. Ann Oncol 2003;14:1660-1666. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/14581275>.

128. Polizzotto MN, Uldrick TS, Wyvill KM, et al. Pomalidomide for symptomatic Kaposi's sarcoma in people with and without HIV infection: a phase I/II study. J Clin Oncol 2016;34:4125-4131. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27863194>.

129. Uldrick TS, Wyvill KM, Kumar P, et al. Phase II study of bevacizumab in patients with HIV-associated Kaposi's sarcoma receiving antiretroviral therapy. J Clin Oncol 2012;30:1476-1483. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22430271>.

130. Schwartzmann G, Sprinz E, Kromfield M, et al. Clinical and pharmacokinetic study of oral etoposide in patients with AIDS-related Kaposi's sarcoma with no prior exposure to cytotoxic therapy. J Clin Oncol 1997;15:2118-2124. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9164226>.

131. Sprinz E, Caldas AP, Mans DR, et al. Fractionated doses of oral etoposide in the treatment of patients with aids-related kaposi sarcoma: a clinical and pharmacologic study to improve therapeutic index. Am J Clin Oncol 2001;24:177-184. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11319295>.

132. Evans SR, Krown SE, Testa MA, et al. Phase II evaluation of low-dose oral etoposide for the treatment of relapsed or progressive AIDS-related Kaposi's sarcoma: an AIDS Clinical Trials Group clinical study. J Clin Oncol 2002;20:3236-3241. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12149296>.

133. Strother RM, Gregory KM, Pastakia SD, et al. Retrospective analysis of the efficacy of gemcitabine for previously treated AIDS-associated Kaposi's sarcoma in western Kenya. *Oncology* 2010;78:5-11. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20215784>.

134. Cao W, Vyboh K, Routy B, et al. Imatinib for highly chemoresistant Kaposi sarcoma in a patient with long-term HIV control: a case report and literature review. *Curr Oncol* 2015;22:e395-399. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26628884>.

135. Koon HB, Bublely GJ, Pantanowitz L, et al. Imatinib-induced regression of AIDS-related Kaposi's sarcoma. *J Clin Oncol* 2005;23:982-989. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15572730>.

136. Koon HB, Krown SE, Lee JY, et al. Phase II trial of imatinib in AIDS-associated Kaposi's sarcoma: AIDS Malignancy Consortium Protocol 042. *J Clin Oncol* 2014;32:402-408. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24378417>.

137. Opravil M, Hirschel B, Bucher HC, Luthy R. A randomized trial of interferon-alpha2a and zidovudine versus bleomycin and zidovudine for AIDS-related Kaposi's sarcoma. *Swiss HIV Cohort Study. Int J STD AIDS* 1999;10:369-375. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10414879>.

138. Shepherd FA, Beaulieu R, Gelmon K, et al. Prospective randomized trial of two dose levels of interferon alfa with zidovudine for the treatment of Kaposi's sarcoma associated with human immunodeficiency virus infection: a Canadian HIV Clinical Trials Network study. *J Clin Oncol* 1998;16:1736-1742. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9586886>.

139. Miles S, Levine A, Feldstein M, et al. Open-label phase I study of combination therapy with zidovudine and interferon-beta in patients with AIDS-related Kaposi's sarcoma: AIDS Clinical Trials Group Protocol 057. *Cytokines Cell Mol Ther* 1998;4:17-23. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9557213>.

140. Fischl MA, Finkelstein DM, He W, et al. A phase II study of recombinant human interferon-alpha 2a and zidovudine in patients with AIDS-related Kaposi's sarcoma. *AIDS Clinical Trials Group. J Acquir Immune Defic Syndr Hum Retrovirol* 1996;11:379-384. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8601224>.

141. Krown SE, Li P, Von Roenn JH, et al. Efficacy of low-dose interferon with antiretroviral therapy in Kaposi's sarcoma: a randomized phase II AIDS clinical trials group study. *J Interferon Cytokine Res* 2002;22:295-303. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12034036>.

142. Krown SE, Lee JY, Lin L, et al. Interferon-alpha2b with protease inhibitor-based antiretroviral therapy in patients with AIDS-associated Kaposi sarcoma: an AIDS malignancy consortium phase I trial. *J Acquir Immune Defic Syndr* 2006;41:149-153. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16394845>.

143. Fortino S, Santoro M, Iuliano E, et al. Treatment of Kaposi's Sarcoma (KS) with nab-paclitaxel. *Ann Oncol* 2016;27:suppl_4: iv124. Available at: <https://doi.org/10.1093/annonc/mdw345.63>.

144. Little RF, Wyvill KM, Pluda JM, et al. Activity of thalidomide in AIDS-related Kaposi's sarcoma. *J Clin Oncol* 2000;18:2593-2602. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10893291>.

145. Fife K, Howard MR, Gracie F, et al. Activity of thalidomide in AIDS-related Kaposi's sarcoma and correlation with HHV8 titre. *Int J STD AIDS* 1998;9:751-755. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9874123>.

146. Nasti G, Errante D, Talamini R, et al. Vinorelbine is an effective and safe drug for AIDS-related Kaposi's sarcoma: results of a phase II study. *J Clin Oncol* 2000;18:1550-1557. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10735904>.